

Table 2: Maternal complications.

	Group 1 (n=316)	Group 2 (n=103)	p value
Cord prolapse	5 (1.6)	1 (0.9)	0.644
Placental abruption	10 (3.2)	5 (4.9)	0.433
Stillbirth	1 (0.3)	1 (0.9)	0.259
Maternal sepsis	1 (0.3)	0 (0.0)	0.564
Maternal fever	33 (10.6)	17 (16.5)	0.112
Postpartum curettage	10 (3.3)	3 (2.9)	0.879
Uterine atony	9 (2.9)	4 (3.9)	0.618
Suspicion of clinical chorioamnionitis	59 (19.0)	15 (14.6)	0.584
Clinical chorioamnionitis	22 (7.1)	7 (6.8)	0.584
Endometritis	3 (1.0)	1 (1.0)	0.996
Blood transfusion	6 (2.0)	1 (1.0)	0.513
Hysterectomy	2 (0.6)	0 (0.0)	0.415
ICU admission	6 (2.0)	0 (0.0)	0.156
Days at hospital after delivery	3 (2-3)	3 (2-3)	0.397

Continuous variables are presented as median (25th - 75th percentile). Categorical variables are presented as number of patients (percentage). Univariate analysis was performed using the Chi-square or Fisher's exact test for comparison of qualitative variables. For quantitative variables, the Student's t test was used for independent samples or the Mann-Whitney U test if the applicability criteria were not met.